

Transplanckian Entanglement Entropy

Darwin Chang¹, Chong-Sun Chu^{1,2} and Feng-Li Lin³

¹ Department of Physics, National Tsing-Hua University, Hsin-Chu, 300 Taiwan

² Department of Mathematics, University of Durham, DH1 3LE, United Kingdom and

³ Physics Division, National Center for Theoretical Sciences,
National Tsing-Hua University, Hsin-Chu, 300 Taiwan

The entanglement entropy of the event horizon is known to be plagued by the UV divergence due to the infinitely blue-shifted near horizon modes. In this paper we calculate the entanglement entropy using the transplanckian dispersion relation, which has been proposed to model the quantum gravity effects. We show that, very generally, the entropy is rendered UV finite due to the suppression of high energy modes effected by the transplanckian dispersion relation.

After 30 years of study, the microscopic origin of the Bekenstein-Hawking entropy [1] of a Schwarzschild black hole remains mysterious. This thermodynamic entropy is finite and is proportional to the area of the event horizon. Matter field contribution to the entropy can be calculated by tracing over the fluctuations beyond the horizon, and is known as the quantum entanglement entropy. The entanglement entropy for black hole was first calculated by 't Hooft [2] by using the so called brick wall method, later on, it was generalized to arbitrary bounded hypersurface [3] and to Rindler space [4, 5]. In [6] a covariant method was developed. In all these results, the entanglement entropy obeys an area law. However, they all suffer from the usual UV divergence of quantum field theory in curved spacetime. This UV divergence is related [4] to the Hawking information paradox.

Intuitively the UV divergence can be understood to be caused by the infinitely blue-shifted near horizon modes. One of the underlying assumptions in these calculations is that the energy-momentum relation takes the same form up to arbitrary high energies. However there are convincing reasons to believe that the conventional picture of spacetime structure, as well as spacetime symmetries, will get modified or even lose its meaning at energy scale comparable to the Planck scale. In a viable theory of quantum gravity, e.g. string theory, there is no UV divergence. It is therefore natural to expect that quantum gravity effects (maybe together with other effects) will suppress the transplanckian modes and hence improve the UV behavior of the entanglement entropy, or even cure the divergences. Although there is so far no proof of it from a theory of quantum gravity, it has been proposed in [8, 10] that one may model the quantum gravity effect on matter propagation by modifying the dispersion relation such that the transplanckian modes are suppressed. In this letter we show that there is a generic class of dispersion relations with such suppression which indeed gives rise UV finite entanglement entropy!

We first review the WKB brick wall method introduced by 't Hooft [2] and apply it to calculate the entanglement entropies for Rindler space, black hole (BH) and the de Sitter space. Then we show that within a general class

of transplanckian dispersion relation (TRD), the entanglement entropy is UV finite. The condition that this class of TRD has to satisfy is shown to be actually very generic. We then give explicit results for the entanglement entropy based on the TRD proposed by Unruh [8] as example. Finally we conclude with a few remarks about the implications of our results.

Entanglement Entropy and UV divergence.— Consider a massless scalar in a 4-dimensional curved spacetime described by the metric $g_{\mu\nu}$ with an event horizon located at r_h . The usual dispersion relation is $g_{\mu\nu}p^\mu p^\nu = 0$. The entanglement entropy was found to be divergent due to the infinitely blue-shifted near horizon region [2, 3]. In the brick wall model, a UV cutoff is introduced to regularize the divergence by imposing the condition on the wavefunction $\Phi(r) = 0$ for $r \leq r_h + \epsilon$, where ϵ is the infinitesimal brick wall thickness [2]. This has the effect of cutting out the near horizon modes.

The total free energy is obtained by summing over the contributions from all the physical modes satisfying the brick wall condition,

$$\beta F = \int d\omega z(\beta\omega) \frac{dg(\omega)}{d\omega} \quad (1)$$

where the Boltzmann weight $z(x) := \ln(1 - e^{-x})$, β is the inverse temperature, and $g(\omega)$ is the density of states. In the WKB approximation [2, 4], it is

$$\pi g(\omega) = \int dr \int \frac{A d^2 p_\perp}{(2\pi)^2} p_r(\omega, \vec{p}_\perp, r), \quad (2)$$

for the Rindler case. Here A is the (regulated) area of the horizon, r and p_r is the radial coordinate and its conjugate momentum, and $p_\perp$ is the transverse momentum. As we will see shortly, the integration limits are determined by the dispersion relation and the above-mentioned brick wall condition. Finally, the entanglement entropy can be obtained from the free energy as $S = \beta^2 \partial F / \partial \beta$.

To be generic, consider a spherically symmetric background metric of the form

$$ds^2 = g_{00} dt^2 + g_{rr} dr^2 + f d\Omega_2^2, \quad (3)$$

where g_{00}, g_{rr} and f are functions of r . The dispersion relation is

$$g^{00}\omega^2 + g^{rr}p_r^2 + p_\perp^2 = 0, \quad (4)$$

where $g^{00} = 1/g_{00}, g^{rr} = 1/g_{rr}$ and $p_\perp^2 = g^{mn}p_m p_n = \ell(\ell+1)/f$ is the transverse momentum squared and ℓ is the angular momentum quantum number. It is convenient to introduce the following definitions

$$\xi^2 := p_\perp^2 g_{rr} = \frac{\ell(\ell+1)g_{rr}}{f}, \quad \rho^2 := -\frac{g_{00}}{g_{rr}}, \quad (5)$$

the dispersion relation (4) takes the form

$$p_r = \sqrt{\omega^2/\rho^2 - \xi^2}. \quad (6)$$

The density of state, $\pi g(\omega) = \int dr d\ell (2\ell+1) p_r$, then takes the form

$$\pi g(\omega) = \int d\rho \mu(\rho) d\xi \xi p_r, \quad (7)$$

where $\mu(\rho)$ is a “measure factor” defined by

$$\mu(\rho) := \frac{2f}{g_{rr}} \frac{dr}{d\rho}. \quad (8)$$

For fixed ω and ρ , we require $\xi \leq \omega/\rho$ to have p_r real. This condition should be imposed in the ξ -integration.

For pedagogical purpose, we now consider some specific cases. First, the Rindler space described by the metric

$$ds^2 = -r^2 dt^2 + dr^2 + dx^2 + dy^2. \quad (9)$$

The event horizon of the local patch is located at $r = 0$. In this case, $\rho = r$, $\xi^2 = p_\perp^2$, $f = A/4\pi$, and the measure factor $\mu(\rho) = A/(2\pi)$. Therefore

$$g(\omega) = \frac{A}{2\pi^2} \int_\epsilon^\infty d\rho \int_0^{\omega/\rho} d\xi \xi p_r = \frac{A\omega^3}{12\pi^2\epsilon^2}. \quad (10)$$

Here a brick wall thickness is introduced in the lower limit of ρ -integration. Together with Eq.(1), we reproduce the result in [4], i.e.,

$$F = -\frac{A\pi^2}{180\beta^4\epsilon^2}, \quad S = \frac{A}{360\pi\epsilon^2}, \quad (11)$$

which are plagued by the UV divergence as $\epsilon \rightarrow 0$. In the evaluation of S , we have set $\beta = 2\pi$ for the inverse Rindler temperature. In [4] the UV divergence was interpreted as an renormalization of the Newton constant.

Using the same method we can also calculate the free energy of the black hole background

$$ds^2 = -h(r)dt^2 + h(r)^{-1}dr^2 + r^2 d\Omega_2^2, \quad (12)$$

where $0 < h(r) = 1 - 2M/r < 1$. In this case $\rho = h$ and $\mu(\rho) = 2(2M)^3\rho/(1-\rho)^4$. The density of state is

$$g(\omega) = \frac{2(2M)^3}{\pi} \int_\epsilon^{1-\delta} d\rho \frac{\rho}{(1-\rho)^4} \int_0^{\omega/\rho} d\xi \xi p_r \quad (13)$$

$$= \frac{2(2M)^3\omega^3}{3\pi} \left(\frac{1}{\epsilon} + \frac{1}{3\delta^3} + \dots \right), \quad (14)$$

where we have introduced an UV (resp. IR) cutoff ϵ (resp. δ) in the ρ -coordinate, which corresponds to an UV (resp. IR) cutoff $r = 2M/(1-\epsilon)$ (resp. $r = 2M/\delta$) in the r -coordinate; and $\dots$ are the sub-leading terms in the limit of $\delta, \epsilon \rightarrow 0$. Using (14) and $\beta = 8\pi M$, we get

$$F = \frac{-2\pi^3(2M)^3}{45\beta^4} \left(\frac{1}{\epsilon} + \frac{1}{3\delta^3} \right), \quad S = \frac{1}{360} \left(\frac{1}{\epsilon} + \frac{1}{3\delta^3} \right) \quad (15)$$

which is plagued by both UV and IR divergences. This agrees with the eqn. (3.12) of [2] except that a IR divergent boundary contribution had been dropped there. The IR-divergent piece is independent of M and represents the contribution from the vacuum. In [2], the UV-divergent piece was shown to give an entropy which obeys the area law $S = \frac{A}{360\pi\epsilon_p^2}$ if the UV-cutoff is given in terms of the proper distance, i.e. $\epsilon_p \approx 4M\sqrt{\epsilon}$. This is the same form as (11). In modern language, the relation between ϵ and M for fixed ϵ_p required by the area law indicates a holographic nature of UV/IR connection [7].

Finally we consider the example of de Sitter space described by the metric (12) but with $0 < h(r) = 1 - r^2/R^2 \leq 1$. In this case, we have again $\rho = h$, but the measure factor is now $\mu(\rho) = R^3\rho\sqrt{1-\rho}$. We find the same result (11) if the UV-cutoff is in terms of $\epsilon_p \approx R\sqrt{\epsilon}$.

In conclusion: The entanglement entropy associated with the event horizon obeys the universal area law, $S = A/(360\pi\epsilon_p^2)$, for the UV cutoff ϵ_p , and diverges as $\epsilon_p \rightarrow 0$.

Transplanckian entropy.— The above UV divergence is due to the infinite contribution of the near horizon modes which have energy far beyond the Planck scale. It is quite generally believed that nonlocal effects due to quantum gravity will provide a natural regulator to the UV divergence in quantum field theory by suppressing the contributions of the high energy modes. It is therefore very natural to expect that in a consistent theory of quantum gravity, the above results will become finite. The simplest proposal to encode the effects of this suppression is to replace the linear dispersion relation by the so-called transplanckian dispersion relation (TDR) [8, 10].

Consider a spherically symmetric background given by the general metric (3), it is reasonable to impose the TDR according to the residual spacetime symmetry preserved by the metric. Since the transplanckian effect will be mostly due to the blue-shift in the near horizon regime in the radial direction, so it is natural to impose TDR along that direction to suppress the blue-shift effect. Therefore we consider TDR of the form

$$g^{00}\omega^2 + g^{rr}H^2(p_r) + p_\perp^2 = 0. \quad (16)$$

The function H describes the transplanckian effects. In terms of the ρ and ξ variables, the TDR (16) then takes the form

$$p_r = H^{-1}(\sqrt{\omega^2/\rho^2 - \xi^2}). \quad (17)$$

For example, Unruh [8] and respectively, Corley and Jacobson [10] (C-J) proposed

$$\text{Unruh : } H_n^U(k) = k_0 [\tanh(k/k_0)^n]^{\frac{1}{n}}, \quad (18)$$

$$\text{C-J : } H^{CJ}(k) = \sqrt{k^2 - k^4/(4k_0^2)}, \quad k \leq 2k_0. \quad (19)$$

The corresponding radial momentum is

$$\text{Unruh : } p_r = k_0 \{ \tanh^{-1} [k_0^{-2}(\omega^2/\rho^2 - \xi^2)]^{\frac{n}{2}} \}^{\frac{1}{n}}, \quad (20)$$

$$\text{C-J : } p_r^{\pm} = \sqrt{2}k_0 \sqrt{1 \pm \sqrt{1 - k_0^{-2}(\omega^2/\rho^2 - \xi^2)}}. \quad (21)$$

Both Unruh's and C-J's proposals belong to a general class of TDR in which the suppression of high energy modes is effected by a bounded function H , says,

$$0 \leq H^2(k) \leq k_0^2, \quad (22)$$

for some k_0 . It follows that for this bounded class of TDR

$$\max(\omega^2/\rho^2 - k_0^2, 0) \leq \xi^2 \leq \omega^2/\rho^2. \quad (23)$$

Combining with the brick wall condition $\rho \geq \epsilon$, we have quite generally the following integration branches contributing to the free energy (up to a factor of $1/(\beta\pi)$)

$$(1) \int_{\epsilon k_0}^{\omega_m} d\omega z \frac{d}{d\omega} \int_{\epsilon}^{\omega/k_0} d\rho \mu(\rho) \int_{\sqrt{\omega^2/\rho^2 - k_0^2}}^{\omega/\rho} d\xi \xi p_r, \quad (24)$$

$$(1*) \int_{(1-\delta)k_0}^{\infty} d\omega z \frac{d}{d\omega} \int_{\epsilon}^{1-\delta} dh \mu(h) \int_{\sqrt{\omega^2/h^2 - k_0^2}}^{\omega/h} d\xi \xi p_r, \quad (25)$$

$$(2) \int_{\epsilon k_0}^{\omega_m} d\omega z \frac{d}{d\omega} \int_{\omega/k_0}^{\rho_m} d\rho \mu(\rho) \int_0^{\omega/\rho} d\xi \xi p_r, \quad (26)$$

$$(3) \int_0^{\epsilon k_0} d\omega z \frac{d}{d\omega} \int_{\epsilon}^{\rho_m} d\rho \mu(\rho) \int_0^{\omega/\rho} d\xi \xi p_r, \quad (27)$$

where $\rho_m = \omega_m = \infty$ for Rindler case, $\rho_m = 1 - \delta$, and $\omega_m = (1 - \delta)k_0$ for BH and de Sitter case. Note that by comparing the ranges of the above ξ -integrations with the ones without transplanckian suppression, we find that the use of TDR leads to a reduction of the allowed angular momentum modes. This leads to a suppression on the density of states and hence the UV finiteness of the entropy as shown below.

It is convenient to change variable $x = \sqrt{\omega^2/\rho^2 - \xi^2}$ in order to isolate the ω -dependence in $g(\omega)$. We have

$$\pi g(\omega) = \int d\rho \mu(\rho) \int dx x H^{-1}(x). \quad (28)$$

Note that the integrand does not depend on ω and ϵ , but the integration limits may. Note also that the (1*)-branch only exists for BH and de Sitter due to the additional constraint $h \leq 1$.

UV Finiteness.— Our goal is to determine the UV behavior of F and S for a generic bounded TDR with a H satisfying (22) such that there are the branches (24)-(27). One may try to estimate the UV behavior for the

integrals for each branches. However this is rather complicated technically. Surprisingly, without knowing explicitly these integrals, one can show that the sum of the branches (1), (1*), (2) and (3), i.e. the free energy F , is completely UV finite!

To demonstrate this, let us first consider (1*) branch. The integration limits are independent of ω , i.e. $\int_{\epsilon}^{1-\delta} d\rho \int_0^{k_0} dx$, so is $g(\omega)$. Therefore (1*) does not contribute. For (1) branch the dx -integral is independent of ω , however, there is ω dependence for $g(\omega)$ coming from the integration limits for $d\rho$ -integral. Using the fundamental theorem of calculus, we can carry out the derivative w.r.t ω in (24) and obtain

$$F_{(1)} = \frac{1}{\pi\beta k_0} \int_0^{k_0} dx x H^{-1}(x) \cdot \int_{\epsilon k_0}^{\omega_m} d\omega z(\beta\omega) \mu\left(\frac{\omega}{k_0}\right). \quad (29)$$

Similarly, we can carry out the derivative w.r.t. ω for branch (2) and (3), and obtain

$$F_{(2)} = -F_{(1)} + \frac{1}{\pi\beta} \int_{\epsilon k_0}^{\omega_m} d\omega z \int_{\frac{\omega}{\rho_m}}^{k_0} dx \mu\left(\frac{\omega}{x}\right) H^{-1}(x), \quad (30)$$

$$F_{(3)} = \frac{1}{\pi\beta} \int_0^{\epsilon k_0} d\omega z \int_{k_0}^{\frac{\omega}{\epsilon}} dx \mu\left(\frac{\omega}{x}\right) H^{-1}(x) + \frac{1}{\pi\beta} \int_0^{\epsilon k_0} d\omega z \int_{\frac{\omega}{\rho_m}}^{k_0} dx \mu\left(\frac{\omega}{x}\right) H^{-1}(x). \quad (31)$$

Physically it is required that the integral

$$I := \int_0^{k_0} dx x H^{-1}(x) < \infty \quad (32)$$

to be finite. The reason is because the density of states for branch (1), in (24), is given by $\pi g(\omega) = I \cdot \int_{\epsilon}^{\omega/k_0} d\rho \mu(\rho)$, and it should be finite for any sensible physical system. This condition also guarantees our manipulation above for (29), (30) to make sense. Of course (32) is true for both Unruh's and C-J's TDRs.

Next we examine the first line of (31), we perform a change of variables by $\omega = \epsilon k_0 y$ and obtain

$$\frac{k_0 \epsilon}{\pi\beta} \int_0^1 dy z(\beta \epsilon k_0 y) \int_{k_0}^{k_0 y} dx \mu\left(\frac{\epsilon k_0 y}{x}\right) H^{-1}(x). \quad (33)$$

Using the fact $z(x) \approx \ln(x) - x/2 + \dots$ around $x = 0$, and assuming the following condition on the near horizon geometry

$$\epsilon \mu(\epsilon) \approx \epsilon^\alpha \quad \text{for } \alpha > 0, \quad \text{as } \epsilon \rightarrow 0, \quad (34)$$

which is true for all three metrics considered, we can manipulate (33) by interchanging the order of integrations over dx and dy so that the y -integration can be carried out, (33) then becomes

$$\frac{\epsilon^\alpha \ln \epsilon}{\pi \alpha k_0} \int_0^{k_0} dx x H^{-1}(x). \quad (35)$$

This vanishes as $\epsilon \rightarrow 0$.

Finally, the second line of (31) can be combined with $F_{(2)}$ of (30), and we obtain for the total free energy

$$F = \frac{1}{\pi\beta} \int_0^{\omega_m} d\omega z(\beta\omega) \int_{\frac{\omega}{\rho_m}}^{k_0} dx \mu\left(\frac{\omega}{x}\right) H^{-1}(x). \quad (36)$$

Note that F , and hence S , is independent of ϵ , i.e. UV finite. This result is quite general and is true as long as (i) the transplanckian modification of the radial modes takes the form (17) with the suppression condition (22), (ii) the metric satisfies a near horizon condition, which expressed in terms of the measure factor as (34). We thus see that TDR generally yields a UV finite free energy and entanglement entropy. This is the main result of this paper.

Examples.— Now we would like to present some more explicit results. We will consider the Unruh dispersion relation with $n = 1$ as an example. For the Rindler case, all the integrals can be worked out explicitly for all three branches. The results are

$$F_{(1)} = \frac{Ak_0^2}{4\pi^2\beta^2} \int_{\beta\epsilon k_0}^{\infty} dy \ln(1 - e^{-y}) \approx -\frac{Ak_0^2}{24\beta^2}, \quad (37)$$

$$F_{(2)} = (2\ln 2 - 1)F_{(1)}, \quad (38)$$

where the approximation in (37) is justified by the infinitesimal $\beta\epsilon k_0$, and

$$F_{(3)} = \frac{Ak_0^3\epsilon}{4\pi^2\beta} \int_0^1 dx \ln(1 - e^{-\beta k_0 \epsilon x}) \left[x \ln \frac{1+x}{1-x} + \ln(1-x^2) \right]. \quad (39)$$

It is easy to show that, with k_0, β fixed, $F_{(3)} \rightarrow 0$, as $\epsilon \rightarrow 0$. Therefore the total free energy and entropy are

$$F = -\frac{\ln 2}{12} \frac{Ak_0^2}{\beta^2}, \quad S = \frac{\ln 2}{12\pi} Ak_0^2. \quad (40)$$

Note that k_0 plays the role of an UV regulator. In fact with $k_0 \rightarrow c/(\beta\epsilon)$, the total free energy is of the form $F = (A/\epsilon^2)\beta^{-4}f(c)$. For $c = 3.4645$, $f(c) = -\pi^2/180$, the result reduces to (11). This is expected since $H_1^U(k) \xrightarrow{k_0 \rightarrow \infty} k$. Moreover, as long as k_0 is far less than the Planck energy scale, the entanglement entropy is just a small correction to the Bekenstein-Hawking entropy.

For the case of BH, the integrals cannot be evaluated for each individual branches. Nevertheless our general result (36) gives the leading IR piece

$$F \approx -\frac{2k_0^3}{3\pi\beta} \left(\frac{2M}{\delta}\right)^3 \int_0^1 dy y \ln(1 - e^{-\beta k_0 y}) \tanh^{-1} y \quad (41)$$

where we assume a large IR cutoff $2M/\delta (\gg \beta)$ to extract the leading IR term from the h -integration. The IR divergence is independent of the background M as in the non-transplanckian case. We note that eqn. (41) is of

the form $F = (2M/\delta)^3 \beta^{-4} g(\beta k_0)$ with $\lim_{\alpha \rightarrow \infty} g(\alpha) = -2\pi^3/135$. Therefore it reproduces the result (15) either for $k_0 \rightarrow \infty$ with β fixed or for the zero temperature limit $\beta \rightarrow \infty$ with k_0 fixed. In general, by performing a numerical study on (41), we find that, for fixed k_0 , both the free energy and the entropy are monotonically decreasing functions of β .

For the case of de Sitter, eqn. (36) gives a result of the form $F = (R^3/\beta^4)j(\beta k_0)$ which is free of both UV and IR divergences. However, $j(x)$ is a complicated function and yields no area law for the entanglement entropy even in the low temperature limit, i.e. $\beta = 2\pi R \rightarrow \infty$. The results for the Corley-Jacobson dispersion relation is similar. We will skip it here.

Discussions.— In this letter we show that the free energy and the entanglement entropy for a wide class of spacetime (so long as the finiteness condition on the “measure factor” is satisfied) are UV finite when the high energy modes are suppressed with the use of TDR. What we show is that for a certain class of TDR, the density of states at high energies does get reduced to a sufficient degree such that the entanglement entropy is UV finite. This is quite a remarkable result because a priori it is not obvious at all. We would like to emphasize that this UV finiteness is largely due to a modification of the propagator, which can be think of as an *UV regulator*, high momentum modes do not need to be *cut out*.

Our results also show that, with the quantum correction, the entanglement entropy may have a complicated area dependence and does not conform to a simple area law in general. This is expected since quantum gravity corrections generally induces higher derivative gravity correction terms that modify the area law, see for examples [11, 12]. It is natural that such modification is also seen in the matter part; and we have shown that this is indeed the case.

We note that without transplanckian suppression, the near horizon modes reproduce the UV behavior of the full integration. However, this is not the case in the transplanckian case. If one keeps only the $\rho = \epsilon$ region in the ρ -integral in evaluating the free energy, we find that the near horizon modes do yield UV improved results, which is one order less divergent than the ordinary case. However it does not render the result finite, at least for the Rindler case, as is given by the full integration. This indicates that the nonlinear suppression feature of the TDR plays a crucial role in improving the UV behavior.

As we mentioned, the TDR of interests are typically characterized by a suppression scale k_0 as in (22). We remark that the transplanckian scale k_0 plays the role of a UV regulator. In fact we have shown for the Rindler case that, by taking $k_0 \sim (\beta\epsilon)^{-1}$, the ordinary UV divergent result is recovered. One can show this similarly for the generic case.

It deserves further study on the thermodynamical behavior of these transplanckian QFT systems and on the

origin of the TDR in the context of string theory, see for example [13] for a discussion. Also it is interesting to re-examine the information paradox based on our result and the earlier study of non-thermal transplanckian Hawking radiation [9, 10].

We thank Bin Chen, Ko Furuta, Pei-Ming Ho, Ted Jacobson and Hoi-Lai Yu for discussions and comments. We acknowledge grants from Nuffield foundation of UK, NCTS and NSC of Taiwan.

-
- [1] J. D. Bekenstein, Phys. Rev. D **7** (1973) 2333;
S. W. Hawking, Commun. Math. Phys. **43** (1975) 199.

- [2] G. 't Hooft, Nucl. Phys. B **256** (1985) 727.
[3] M. Srednicki, Phys. Rev. Lett. **71** (1993) 666.
[4] L. Susskind, J. Uglum, Phys. Rev. D **50** (1994) 2700.
[5] C. G. Callan, F. Wilczek, Phys. Lett. B **333** (1994) 55.
[6] S. N. Solodukhin, Phys. Rev. D **51** (1995) 609.
[7] L. Susskind, E. Witten, hep-th/9805114.
[8] W. G. Unruh, Phys. Rev. D **51** (1995) 2827.
[9] R. Brout, S. Massar, R. Parentani, Ph. Spindel Phys. Rev. **D52** (1995) 4559.
[10] S. Corley, T. Jacobson, Phys. Rev. D **54** (1996) 1568.
[11] G. L. Cardoso, B. de Wit, T. Mohaupt, Phys. Lett. **B451** (1999) 309; Nucl. Phys. **B567** (2000) 87.
[12] S. Carlip, Class. Quant. Grav. **17** (2000) 4175.
[13] C. S. Chu, B. R. Greene, G. Shiu, Mod. Phys. Lett. A **16** (2001) 2231.